HYPROMELLOSE 0.3% W/V EYE DROPS

for you. Ask your pharmacist if you need any more information or advice.	5ml,10ml,15ml & 20ml of solution. Not all pack sizes may be marketed.
(PRIOPILOS Hypromellose Eye Drops is a lubricating artificial tear drops used to relieve dry eye conditions. It provides a smooth feeling for sore eyes with soothing relief from grittiness, tichniess, buming and freeigh obby exercision. WHAT CAUSES BRY EVE CONDITION Dry eye condition associated with the use of VDUs and TVs, infrequent blinking, certain medical treatments, atmospheric pollution and drying atmospheres eg, air-conditioning, certain healting, wind and sur it is also result of ocular surface disorders that can be caused by a number of different reasons where the normal teas are either of power quite promoted in the certain surface of the production of the certain of the cer	INSTRUCTION FOR USE First wesh your hands Tighten the cap on the nozile to pierce the tip of the eye drop bottle. Toghten the cap on the nozile to pierce the seal and remove the cap. If you are a contact lens wearer remove your lenses and do not replace until 30 minutes after using eye drops. Till your head back and look at the ceiling. Pull the bover eyelf centryl downwards. Hold the bottle upsade down above the eye and gently squeeze the bottle to release a drop into your eye. Aeried touching any part of the eye (or any other surface) with the tip of the bottle. The closed eye, and hold for 1 minute of the chapter of the closed eye, and hold for 1 minute. Repals for the their eye if necessite of the with a picked cap. When using the bottle for the first time, screw the cap down tightly in order to pierce the tip of the bottle.
The usual dosage is one or two drops to be applied 3 to 4 times a day as needed, or as directed by your doctor. No dosage adjustment is required in children	
WARNINGS AND PRECAUTIONS	
Talk to your doctor or pharmacist before using Hypromellose 0.3% w/v Eye Drops	
If you are wearing contact lenses. If you are wearing contact lenses. If you are allergic to Hypromellose or any other ingredient of this eye drops	Eye solutions, if handled wrongly, can become contaminated by common bacteriand cause eye infections.
Other Medicines And Hypromellose	Distributed by:
Please tell your doctor or pharmacist if you are applying any other type of eye drops or eye ointment before you start to use this eye drops	Biovantic Pharma Limited Biovantic Kemp House, 160 City Road, London, EC1V 2NX, UK.
Pregnancy and Breast-feeding	
If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this eye drops	
Driving and Using Machines	
Hypromellose may cause a temporary blurring of vision immediately after using the drops. Do not drive or operate machines if you experience any visual disturbance after using the product. Wait until this clears before driving or using machines.	
POSSIBLE SIDE EFFECTS	
Possible side effects include: - Blurred vision - Eve pain	
- Foreign body sensation in eyes - Eye irritation - Redness of eye	A STERILEA C€ II ⊗ ⊗ 1 30°C 35°C
These effects should wear off after a short time. If these effects persist or get worse	⚠ Warning
stop using Hypromellose and talk to your doctor or pharmacist. COMPOSITION:	LOT Lot No.
Each ml Contains:	∑ Use By
Hypromellose 0.3% w/v	Manufacturer
Benzalkonium Chloride Solution 0.02% w/v (As Preservative) The other ingredients are Sodium Chloride, Potassium Chloride, Borax, Boric Acid,	STERILE A Symbol for Sterile medical devices processed using assotic techniques
Sodium Hydroxide Solution, Hydrochloric Acid STORAGE	Please follow the instructions for use
Keep this eye drops out of the sight and reach of children. Do not use this eye drops after the expiry date. Do not share eye drops with any other person.	The product satisfies the basic requirements as set forth in Appendix I of Directive 93/42/EEC concerning medical devices.
Do not store above 30°C. Do not freeze. Protect from light.	Not Injectable
FOR EXTERNAL USE ONLY	Do not use if the vial is damaged.
NOT FOR INJECTION	Keep away from sunlight
Discard the bottle 28 days after opening, even if the solution is remaining.	Date of Issue : Jun 2020 Revision : 03 Revision Date : March 2021